

Influenza Diagnostics Market Trends Overview by Share, Size, Growth and Competitive Landscape 2022-2030

Influenza Diagnostics Market Trends Overview by Share, Size, Growth and Competitive landscape 2022-2030

Market Overview

Market Research Future (MRFR) has published a report about the global influenza diagnostics market share expects 5.3% CAGR during the forecast period between 2022-2030. In terms of cash, this market is expected to grow with USD 1.63 billion by 2030. This report analyzes the market, potential for growth during the forecast period, market sizes around the world, key players in the market, product launches and latest research and development (R&D).

Influenza as an illness can cause severe complications and mess up anybody's medical history. Influenza is of various types and with a variety of deadly symptoms. Hence, early diagnosis of influenza can reduce the chances of further complications. Therefore, with World Health Organization (WHO) confirming about 600 million Influenza cases worldwide, the services offering accurate Influenza diagnosis are going to be in-demand always and become a part of a huge market in the medical sector.

Key Players

Global influenza diagnostics market players include Alere (USA), BD (USA), Becton Dickinson (USA), BioMérieux SA (France), Cepheid (USA), Hardy Diagnostics (USA), LifeSign LLC (USA), Meridian Bioscience, Inc. (USA), Quidel Corporation (USA), Roche Molecular Systems Inc. (Switzerland), and Sekisui Diagnostics (USA).

Market Segmentation

The global [influenza diagnostics market trends](#) is segmented on the basis of types, tests, and region. The segmentation of types comprises of type A flu, type B flu, type C flu. The three forms of type A are bird flu, seasonal flu, and Swine flu. Bird Flu is known as Avian Influenza which is caused by viruses adapted to birds. It can be caused by eating infected birds. Seasonal flu is common cold whose viruses become active only during the change of seasons as human body becomes adapt to the new season slowly. Similar to bird flu, swine flu is an infection caused by any one of several types of swine influenza viruses. Swine influenza virus (SIV) or swine-origin influenza virus (S-OIV) are the viruses causing this infection. These viruses are found in pigs. Type B flu is found only in humans. Influenza type B viruses are not classified by subtype and do not cause pandemics. It is caused by a genus called Influenza virus B which is from the virus family Orthomyxoviridae. Type C Influenza viruses are from Orthomyxoviridae family. They are not as severe as Type A or B but can cause local epidemics.

On the basis of tests, the market has been segmented into direct fluorescent antibody (DFA) tests, molecular tests, nucleic acid sequence-based amplification (NASBA) tests, rapid influenza detection tests (RIDT), serological assays, simple amplification-based assays (SAMBA) and loop-mediated isothermal amplification-based assays (LAMP). RIDT is the most common test for influenza, as it can confirm influenza in just 30 minutes.

Serological assays are segmented into primary serological tests, secondary serological tests, and tertiary serological tests. Primary serological tests cover enzyme-linked immunosorbent assay (ELISA), immunofluorescent antibody technique (IFAT) and radioimmunoassay (RIA). Secondary Serological tests include agglutination tests, complement fixation tests (CFT), precipitation tests, serum

